

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Keep away from children, pets, and domestic animals. Do not contaminate feed and food stuffs. Use only in rodent-infested areas where exposure to agricultural crops or commodities will not occur. Wash thoroughly with soap and water after handling and before eating or smoking. Clean clothing should be used daily.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds, and other wildlife. Bait exposed to soil surfaces may be hazardous to birds and other wildlife. Do not apply directly to lakes, streams, or ponds.

**STORAGE AND DISPOSAL**

Store in a locked enclosure (in a properly posted area) in such a manner so as not to present a hazard to persons, animals food, feed, crops, or property. Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Completely empty bag into application equipment. Then, dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**RODENT BAIT  
CHLOROPHACINONE  
TREATED GRAIN (0.01%)**

Active Ingredient: Chlorophacinone 2-[(p-chlorophenyl) phenyl-acetyl] 1, 3-indandione:	0.01%
Inert Ingredients:	<u>99.99%</u>
Total:	100.00%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED: CALL A PHYSICIAN OR POISON CONTROL CENTER IMMEDIATELY:** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat. Do not induce vomiting or give anything by mouth to an unconscious person.

**NOTE TO PHYSICIAN**

Intramuscular and oral administration of Vitamin K<sub>1</sub>, combined with blood transfusions is indicated as in the case of hemorrhage caused by overdoses of bis-hydroxycoumarin (Dicumarol).

REGISTRANT: California Department of Food and Agriculture  
1220 "N" Street, Room A357  
Sacramento, CA 94271-0001

CALIF. REG. NO. 10965-50006  
EPA SLN. NO CA 890024  
EPA EST. NO.

**NET CONTENTS:** \_\_\_\_\_

April 94

**DIRECTIONS FOR USE**

It is a violation of State and Federal laws to use this product inconsistent with its labeling.

**FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA  
USE RESTRICTIONS**

For use against GROUND SQUIRRELS (*Spermophilus beecheyi*, *S. beldingi*), POCKET GOPHERS (*Thomomys* spp.), DEER MICE (*Peromyscus* spp.) and HOUSE MICE (*Mus musculus*).

**IMPORTANT**

Bait must be eaten at several feedings on five or more successive days with no period longer than 48 hours between feedings. Keep this product in a closed container, free from odors which may reduce acceptability.

**BAITING:** GROUND SQUIRRELS, *Spermophilus beecheyi*, *S. beldingi*.

Scatter a handful of bait (about 10 baits per pound) evenly over 40 to 50 square feet near active burrows or runways. Repeat every other day for 3 to 4 applications. Do not pile bait.

**BAITING:** DEER MICE, *Peromyscus* spp.

Spread bait evenly by hand, mechanical spreader, or aircraft at the rate of 2 to 6 pounds per acre through the infested area. Sufficient bait should be applied to last a minimum of 4 days (preferably without rain). In reforestation programs bait application should precede the sowing of conifer seed by 10 days to 2 weeks.

**BAITING:** HOUSE MICE, *Mus musculus*.

Place tablespoon (1/4 to 1/2 ounce) of bait in bait box or shallow container, preferably in protected feeder stations. Place bait stations at 8 to 12 foot intervals in dry locations such as in concealed places, in corners, or along walls where house mice feed, drink, or frequent. Inspect stations daily and add bait as needed. Replace moldy or old bait with fresh bait. Dispose of baits in such a manner as to prevent ingestion by children, pets, wildlife, and domestic animals. An uninterrupted supply of bait should be maintained as long as any bait is taken, which may be 2 to 4 weeks. Where continuous source of infestation is present, permanent bait stations should be established and the bait replenished as needed. All poison baits must be picked up or disposed of after rodent activity has stopped. An uninterrupted supply of bait should be available for 6 to 8 days.

For mechanical spreader applications apply at the rate of 10 pounds per acre through infested area, follow with second application in 2 to 3 days.

**BAITING:** POCKET GOPHERS, *Thomomys* spp.

A hand probe should be used to locate main burrow system of gophers. Carefully rotate probe to enlarge opening. Using a funnel, place 1/2 cup of treated bait into burrow. Cover hole with clod or dirt. Do not disturb bait. Bait should be placed in two locations per gopher system. Repeat treatment as necessary. Additional information is available from the agricultural commissioner's office.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Keep away from children, pets, and domestic animals. Do not contaminate feed and food stuffs. Use only in rodent-infested areas where exposure to agricultural crops or commodities will not occur. Wash thoroughly with soap and water after handling and before eating or smoking. Clean clothing should be used daily.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds, and other wildlife. Bait exposed to soil surfaces may be hazardous to birds and other wildlife. Do not apply directly to lakes, streams, or ponds.

**STORAGE AND DISPOSAL**

Store in a locked enclosure (in a properly posted area) in such a manner so as not to present a hazard to persons, animals, food, feed, crops, or property. Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**DIRECTIONS FOR USE CONTINUED**

**BAITING:** JACK RABBITS (Black-tailed hare) *Lepus californicus*.

Place 1 to 5 pounds of bait in a covered self-dispensing feeder or nursery flat in areas frequented by rabbits (near runways, resting or feeding areas). It may be necessary to move feeders to new locations to achieve bait acceptance. Carcasses of poisoned rabbits should be picked up and disposed of by deep burying or burning.

**BAITING:** MEADOW MICE, *Microtus californicus*, *Microtus montanus*.

Spot baiting: Lightly scatter tablespoon amounts (¼ to ½ ounce) of bait near active burrows or in runways. Repeat treatment every other day for three treatments.

**RODENT BAIT  
CHLOROPHACINONE  
TREATED GRAIN (0.005%)**

Active Ingredients: Chlorophacinone 2-[p-chlorophenyl phenyl-acetyl]	
1, 3-indandione	0.005%
Inert Ingredients:	99.995%
Total:	100.000%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED: CALL A PHYSICIAN OR POISON CONTROL CENTER IMMEDIATELY.** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat. Do not induce vomiting or give anything by mouth to an unconscious person.

**NOTE TO PHYSICIAN**

Intramuscular and oral administration of Vitamin K<sub>1</sub>, combined with blood transfusions is indicated as in the case of hemorrhage caused by overdoses of bis-hydroxycoumarin (Dicumarol).

REGISTRANT: California Department of Food and Agriculture  
1220 "N" Street, Room A357  
Sacramento, CA 95814

CALIF. REG. NO. 10965-50004  
EPA SLN. NO. CA 890023  
EPA EST. NO.

**NET CONTENTS:** \_\_\_\_\_

**DIRECTIONS FOR USE**

It is a violation of State and Federal laws to use this product inconsistent with its labeling.

**FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA**

**USE RESTRICTIONS**

For use against NORWAY RATS, ROOF RATS, GROUND SQUIRRELS, CHIPMUNKS, MUSKRATS, JACKRABBITS, MEADOW MICE, WOOD RATS, and GOLDEN-MANTLED GROUND SQUIRRELS.

**IMPORTANT**

An uninterrupted supply of bait should be maintained as long as any bait is taken, which may be 1 to 4 weeks. Treated baits shall be placed in locations not accessible to children, pets, wildlife, and domestic animals, or in tamper-resistant bait boxes. Initial acceptance may not occur until animals become accustomed to the bait station. Bait stations should have entrance holes large enough to admit target species, but not larger animals. Secure bait stations to prevent overturning. Maintain bait supply for at least 15 days. Continue baiting until all signs of feeding have been stopped. Bait placements should be checked daily; replace consumed, contaminated, or spoiled baits. Where a continuous source of infestation is present, permanent bait stations should be established and the bait replenished as needed. All poison baits must be picked up and disposed of after rodent activity has stopped. Keep this product in a closed container to maintain acceptability.

**BAITING:** GROUND SQUIRRELS, *Spermophilus beecheyi*, *S. beldingi*.

Bait boxes: Place 1 to 5 pounds of bait in a covered bait box in areas frequented by ground squirrels (near runways, burrows, etc.)

**BAITING:** GOLDEN-MANTLED GROUND SQUIRRELS, *Spermophilus lateralis*, and CHIPMUNKS, *Eutamias* spp.

Bait boxes: Place 1 to 5 pounds of bait in a covered bait box in areas frequented by target animals (near runways, burrows, etc.).

**BAITING:** MUSKRATS, *Ondatra zibethica*.

Floating bait station: Place 1 to 5 pounds of bait in floating bait box.

**BAITING:** NORWAY RATS, *Rattus norvegicus*, and ROOF RATS, *R. rattus*.

Place 4 to 16 ounces of bait in bait box or shallow container, preferably in protected feeder stations. Place bait stations in dry locations such as in concealed places, in corners, along walls where rats feed, drink or frequent. For roof rats, place bait at ground floor and top floor, or attic levels. For Norway rats, put bait at or near ground level and at burrows and harborages.

**BAITING:** WOOD RATS, *Neotoma* spp. Place 4 to 6 ounces of bait in an open bait box or shallow container. Bait stations should be located near existing wood rat runways or dens, and generally spaced no further than 100 feet apart.

**PRECAUTIONARY STATEMENTS**  
**Hazard to Humans and Domestic Animals**  
**CAUTION**

Keep away from children, pets, and domestic animals. Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Do not contaminate food or food stuffs. Use only in rodent infested areas where exposure to agricultural crops or commodities will not occur. Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. All handlers, including loaders and applicators, must wear: long sleeve shirts and long pants, shoes plus sock, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Keep and wash PPE separate from other laundry. Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water.

**Environmental Hazards**

This product is toxic to fish, birds, and other wildlife. Bait exposed to soil surfaces may be hazardous to birds and other wildlife. Dogs and other predatory and scavenging animals might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high-water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

**ENDANGERED SPECIES  
CONSIDERATIONS**

Notice: Killing of an endangered species may result in fine and/or imprisonment under the Endangered Species Act. The use of this product may pose a hazard to a federally designated endangered/threatened species. For protection of federally listed species, users shall consult the U.S. EPA Endangered Species Bulletin for the county in which the application will occur. A copy of the bulletin may be obtained from the county agricultural commissioner or downloaded from the internet at the following website: <http://www.cdpr.ca.gov/docs/es/colist.htm>.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

Nonrefillable container. Do not reuse or refill this container.

**Pesticide Storage:** Store in a dry place. Do not store around the home. Store in a locked enclosure or other location not accessible to children, pets, domestic animals or wildlife.

**Pesticide Disposal:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal:** When finished with empty containers, dispose of them in a sanitary landfill, or by incineration, or if allowed by State or Local authorities, by burning. If burned, stay out smoke.

**RESTRICTED USE PESTICIDE**

Due to Hazard to Nontarget Organisms  
 For retail sale to and use only by Certified Applicators or persons  
 under their direct supervision and only for those uses covered  
 by the Certified Applicator's certification.

**RODENT BAIT**  
**CHLOROPHACINONE**  
**TREATED ARTICHOKE**  
**BRACTS (0.01%)**

Active Ingredient:  
 Chlorophacinone (2-[(p-Chlorophenyl)phenyl acetyl]-1,3-indandione).....0.01%  
 Other Ingredients.....99.99%  
 Total.....100.00%

**CAUTION**

**FIRST AID**

**Anticoagulant (Bis-hydroxycoumarin class)**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

<b>If Swallowed</b>	<ul style="list-style-type: none"> <li>▪ Call a poison control center or doctor immediately for treatment advice.</li> <li>▪ Have person sip a glass of water if able to swallow.</li> <li>▪ Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>▪ Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on Skin</b>	<ul style="list-style-type: none"> <li>▪ Take off contaminated clothing.</li> <li>▪ Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>▪ Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If Inhaled</b>	<ul style="list-style-type: none"> <li>▪ Move person to fresh air</li> <li>▪ If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.</li> <li>▪ Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If in Eyes</b>	<ul style="list-style-type: none"> <li>▪ Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>▪ Call a poison control center or doctor for treatment advice.</li> </ul>

For more information on this pesticide product (including health concerns and medical emergencies), call the National Pesticide Telecommunications Network at 1-800-858-7378.

**NOTE TO PHYSICIAN**

This product contains chlorophacinone, an anticoagulant of the bis-hydroxycoumarin class. Vitamin K is antidotal.

**NOTE TO VETERINARIAN**

Contains chlorophacinone, an anticoagulant with a half-life in the dog of unknown days. For dogs that have ingested or that are suspected of having ingested chlorophacinone, and/or have obvious poisoning symptoms, such as bleeding or have lowered prothrombin times, give Vitamin K as follows: intramuscular injection. For anticoagulants with long half-lives, if known, it might be necessary to check prothrombin times every 3 days until values return to normal. See 'Note to Physician' for additional information.

**KEEP OUT OF REACH OF CHILDREN**  
**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**THIS PRODUCT MAY ONLY BE DISTRIBUTED AND USED IN CALIFORNIA AND MAY ONLY BE SOLD BY OFFICES OF CALIFORNIA'S COUNTY AGRICULTURAL COMMISSIONERS**

**READ THIS LABEL:** Read this entire label and follow all use directions and precautions.

**IMPORTANT:** Store product not in use in a location out of reach of children, pets, domestic animals, and wildlife. Dispose of product container, unused, spoiled, and unconsumed bait as specified in the "Storage and Disposal" section of this label.

**USE RESTRICTIONS:** This product may only be used to control California Voles (*Microtus californicus*) in artichoke fields in California. Contact your County Agricultural Commissioner's office if you need help in determining whether California voles are damaging your artichokes.

**Do not** apply this product in or around homes or other human residences. **Do not** apply this bait anywhere but to bare ground in vole invested artichoke fields. **Do not** apply this product by any other method other than the method specified in this label.

**Do not** apply this product in occupied habitat of endangered species that may be harmed by exposure to this bait or to animals that have consumed it. Contact your County Agricultural Commissioner's office or <http://www.cdpr.ca.gov/docs/es/colist.htm> for information on endangered species that use of this product in your area might impact.

**Do not** apply this product in any way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in area during application. Keep all other persons out of treatment area during application.

**Do not** apply bait when rain is likely within 24 hours. **Do not** apply on roads or in water or in areas accessible to farm or ranch animals.

**Baiting:** Scatter about one ounce (4-5 bracts) of bait per artichoke plant on bare ground in vole-infested areas. Apply bait near vole burrow openings or runways. Do not overbait! Be careful to scatter bait so that it does not directly contact any parts of the artichoke plant. If necessary, make second and third applications at 21-day intervals. Do not exceed three applications in one 12-month period.

**\*This 24 C SLN will expire on April 30, 2009.**

California Department of Food and Agriculture  
 1220 "N" Street, Room 341  
 Sacramento, CA 95814

EPA SLN No. CA 930022  
 EPA EST. No. XXXX-CA-X  
 California Reg. No. 10965-50067

**Net Contents:**

Revised 3/27/08