

## PRECAUTIONARY STATEMENTS

### Hazard to Humans and Domestic Animals

# CAUTION

Keep away from children, pets, and domestic animals. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear long-sleeved shirts and long pants, socks and shoes, and gloves. Wash thoroughly with soap and water after handling. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Any person who retrieves carcasses or unused bait following application of this product must wear gloves. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove personal protective equipment (PPE) immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Keep and wash PPE separate from other laundry. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water.

### Environmental Hazards

This product is toxic to fish, birds, and other wildlife. Bait exposed to soil surfaces may be hazardous to birds and other wildlife. Dogs and other predatory and scavenging animals might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high-water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

### ENDANGERED SPECIES CONSIDERATIONS

Notice: Killing of an endangered species may result in fine and/or imprisonment under the Endangered Species Act. The use of this product may pose a hazard to a Federally designated endangered/threatened species. For protection of federally listed species, users shall consult the U.S. EPA Endangered Species Bulletin for the county in which the application will occur. A copy of the bulletin may be obtained from the county agricultural commissioner or downloaded from the internet at the following website: <http://www.cdpr.ca.gov/docs/es/colist.htm>.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

**Pesticide Storage:** Store in a dry place. Do not store around the home. Store in a locked enclosure or other location not accessible to children, pets, domestic animals or wildlife.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Handling:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or if allowed by State or Local authorities, by burning. If burned, stay out of smoke.

### RESTRICTED USE PESTICIDE

#### Due to Hazards to Nontarget Organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

## RODENT BAIT DIPHACINONE TREATED GRAIN (0.01%)

Active Ingredient:  
Diphacinone 2-diphenylacetyl-1, 3-indandione: .....0.01%  
Other Ingredients..... 99.99%  
Total:.....100.00%

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

#### FIRST AID

##### Anticoagulant (Bis-hydroxycoumarin class)

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

|                     |  |
|---------------------|--|
| <b>If Swallowed</b> | <ul style="list-style-type: none"> <li>▪ Call a poison control center or doctor immediately for treatment advice.</li> <li>▪ Have person sip a glass of water if able to swallow.</li> <li>▪ Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>▪ Do not give anything by mouth to an unconscious person.</li> </ul> |
| <b>If on Skin</b>   | <ul style="list-style-type: none"> <li>▪ Take off contaminated clothing.</li> <li>▪ Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>▪ Call a poison control center or doctor for treatment advice.</li> </ul>  |
| <b>If in Eyes</b>   | <ul style="list-style-type: none"> <li>▪ Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>▪ Call a poison control center or doctor for treatment advice.</li> </ul>   |

For more information on this pesticide product (including health concerns and medical emergencies), call the National Pesticide Telecommunications Network at 1-800-858-7378.

##### NOTE TO PHYSICIAN

This product contains diphacinone, an anticoagulant of the bis-hydroxycoumarin class. Vitamin K is antidotal. If ingested, administer Vitamin K intramuscularly or orally, as indicated in bis-hydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

##### NOTE TO VETERINARIAN

If a pet or other unintended animal ingests the bait call a veterinarian or 1-800-858-7378 at once. Contains diphacinone, an anticoagulant with a half-life in the dog of unknown days. For dogs that have ingested or that are suspected of having ingested diphacinone, and/or have obvious poisoning symptoms, such as bleeding or have elevated prothrombin times, give Vitamin K as follows: intramuscular injection. For anticoagulants with long half-lives, if known, it might be necessary to check prothrombin times every 3 days until values return to normal. See 'Note to Physician' for additional information.

California Department of Food and Agriculture  
1220 "N" Street, Room 341  
Sacramento, CA 95814

EPA SLN No. CA 890022  
California Reg. No. 10965-50003

EPA EST. No. XXXX-CA-X  
Net Contents:

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**THIS PRODUCT MAY ONLY BE DISTRIBUTED AND USED IN CALIFORNIA AND MAY ONLY BE SOLD BY THE OFFICES OF CALIFORNIA'S COUNTY AGRICULTURAL COMMISSIONERS.**

**READ THIS LABEL:** Read this entire label and follow all use directions and precautions.

**IMPORTANT:** Store product not in use in a location out of reach of children, pets, domestic animals, and wildlife. Dispose of product container, unused, spoiled, or unconsumed bait as specified in the "STORAGE AND DISPOSAL" section of this label.

#### USE RESTRICTIONS – FOR ALL USES

This product may only be used to control CALIFORNIA GROUND SQUIRRELS (*Spermophilus beecheyi*), DEER MICE (*Peromyscus spp.*) and HOUSE MICE (*Mus musculus*), at the use sites and using the application methods identified in the "USE RESTRICTIONS:" paragraphs indicated below for the target species groupings. Contact your local County Agricultural Commissioner's office if you need help in identifying the target species you intend to control or are not sure about the limits of the site designations.

Use only in rodent infested areas where exposure to agricultural crops or commodities will not occur. Do not contaminate food or food stuffs.

Do not apply this product in or around homes or other human residences. Do not apply this bait at sites or to control pests not indicated on this label. Do not apply this product by application methods not specified in this label. Do not pile bait.

Do not apply this product in occupied habitat of endangered species that may be harmed by exposure to this bait or to animals that have consumed it. Contact your local County Agricultural Commissioner's office or <http://www.cdpr.ca.gov/docs/endspec/prescint.htm> for information on endangered species that use of this product in your area might impact.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Keep all other persons out of the treatment area during application.

Do not apply bait in or near actively growing food or feed crops or in areas where contaminating food or surfaces that may contact food is likely. Do not apply bait by aircraft at sites where bait may contaminate bodies of water. When applying bait along canals and other waterways, use equipment that directs bait away from the water.

#### FOLLOW-UP OPERATIONS – FOR ALL USES

Collect dead rodents and dispose of them by deep burying, burning (if permitted in your County or community), or by double-plastic bagging or wrapping in newspaper and discarding in trash. Wear disposable plastic gloves or other suitable hand protection if you must pick up carcasses by hand.

#### **VINEYARDS, ORCHARDS, GROVES, FORESTRY PLANTATIONS, PASTURES, RANGELANDS AND SELECTED NONCROP AREAS**

#### CALIFORNIA GROUND SQUIRRELS

**USE RESTRICTIONS:** This product may be used to control California Ground Squirrels (*Spermophilus beecheyi*) in broadcast bait applications in vineyards, orchards and groves (non-bearing season only); non-crop borders, and fallow lands; along outsid es of fence rows and rights-of-way adjacent to canal banks, ditch banks, highways, levees, railroad lines, and utilities; and in campgrounds, recreational areas, horticultural nurseries, plantations of forest trees, pastures and rangelands.

Do Not graze livestock or plant food or feed crops in spot-treated areas while bait is present. Applications in orchards, groves, and vineyards may only be made after harvest and during the dormant period and may not be made after tree and vine growth resumes in the spring.

**BAIT ACCEPTANCE TRIALS:** Before applying toxic bait, test for target species' readiness to accept this product by manually scattering untreated crimped oat groats to portions of the infested area. Using flagging or other suitable items, mark areas where oats were scattered and return on the following day to observe where oats are being consumed to assess whether consumption is by ground squirrels or nontarget species. Do not apply toxic bait if it appears that nontarget species are primarily responsible for consuming the oats. Apply toxic bait only if ground squirrels appear to have accepted the oats readily. Untreated crimped oat groats may be obtained from the County Agricultural Commissioner's office.

**BROADCAST BAITING:** Using a ground-based mechanical spreader or aircraft, apply bait at 10 pounds per swath acre, depending upon the density of ground squirrels. Make a second application 4 days after the first treatment.

#### **DEER MICE**

**USE RESTRICTIONS:** This product may be used to control deer mice (*Peromyscus spp.*) in vineyards, orchards and groves (non-bearing season only), non-crop borders, and fallow lands; along outsides of fence rows and rights-of-way adjacent to canal banks, ditch banks, highways, levees, railroad lines, and utilities; and in campgrounds, recreational areas, horticultural nurseries, plantations of forest trees, pastures and rangeland.

**Do Not** graze livestock or plant food or feed crops in spot-treated areas while bait is present. Applications in orchards, groves, and vineyards may only be made after harvest and during the dormant period and may not be made after tree and vine growth resumes in the spring.

**BROADCAST BAITING:** Using a ground-based mechanical spreader or aircraft, apply bait at 2 to 6 pounds per swath acre, depending upon the density of deer mice. Make a second application 4 days after the first treatment.

#### **HOUSE MICE**

**USE RESTRICTIONS:** This product may be used to control house mice (*Mus musculus*) in vineyards, orchards and groves (non-bearing season only), non-crop borders, and fallow lands; along outsides of fence rows and rights-of-way adjacent to canal banks, ditch banks, highways, levees, railroad lines, and utilities; and in campgrounds, recreational areas, horticultural nurseries, plantations of forest trees, pastures and rangelands.

**Do Not** graze livestock or plant food or feed crops in spot-treated areas while bait is present. Applications in orchards, groves, and vineyards may only be made after harvest and during the dormant period and may not be made after tree and vine growth resumes in the spring.

**BROADCAST BAITING:** Using a ground-based mechanical spreader or aircraft, apply bait at 6-10 pounds per swath acre, depending upon the density of house mice. Make a second application 4 days after the first treatment.

Revised 2/24/11

## PRECAUTIONARY STATEMENTS

### Hazard to Humans and Domestic Animals

## CAUTION

Keep away from children, pets, and domestic animals. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear long-sleeved shirts and long pants, socks and shoes, and gloves. Wash thoroughly with soap and water after handling. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Any person who retrieves carcasses or unused bait following application of this product must wear gloves. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove personal protective equipment (PPE) immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Keep and wash PPE separate from other laundry. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water.

### Environmental Hazards

This product is toxic to fish, birds, and other wildlife. Bait exposed to soil surfaces may be hazardous to birds and other wildlife. Dogs and other predatory and scavenging animals might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high-water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

## ENDANGERED SPECIES CONSIDERATIONS

Notice: Killing of an endangered species may result in fine and/or imprisonment under the Endangered Species Act. The use of this product may pose a hazard to a Federally designated endangered/threatened species. For protection of federally listed species, users shall consult the U.S. EPA Endangered Species Bulletin for the county in which the application will occur. A copy of the bulletin may be obtained from the county agricultural commissioner or downloaded from the internet at the following website: <http://www.cdpr.ca.gov/docs/es/colist.htm>.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

**Pesticide Storage:** Store in a dry place. Do not store around the home. Store in a locked enclosure or other location not accessible to children, pets, domestic animals or wildlife.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Handling:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or if allowed by State or Local authorities, by burning. If burned, stay out of smoke.

## RESTRICTED USE PESTICIDE

### Due to Hazard to Nontarget Organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

## RODENT BAIT DIPHACINONE TREATED GRAIN (0.005%)

Active Ingredient:  
Diphacinone 2-diphenylacetyl-1, 3-indandione:.....0.005%  
Inert Ingredients..... 99.995%  
Total:.....100.000%

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

#### FIRST AID

##### Anticoagulant (Bis-hydroxycoumarin class)

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

|                     |   |
|---------------------|---|
| <b>If Swallowed</b> | <ul style="list-style-type: none"><li>Call a poison control center or doctor immediately for treatment advice.</li><li>Have person sip a glass of water if able to swallow.</li><li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>Do not give anything by mouth to an unconscious person.</li></ul> |
| <b>If on Skin</b>   | <ul style="list-style-type: none"><li>Take off contaminated clothing.</li><li>Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>Call a poison control center or doctor for treatment advice.</li></ul>   |
| <b>If in Eyes</b>   | <ul style="list-style-type: none"><li>Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>Call a poison control center or doctor for treatment advice.</li></ul>   |

For more information on this pesticide product (including health concerns and medical emergencies), call the National Pesticide Telecommunications Network at 1-800-858-7378.

#### NOTE TO PHYSICIAN

This product contains diphacinone, an anticoagulant of the bis-hydroxycoumarin class. Vitamin K is antidotal. If ingested, administer Vitamin K intramuscularly or orally, as indicated in bis-hydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

#### NOTE TO VETERINARIAN

If a pet or other unintended animal ingests bait call a veterinarian or 1-800-858-7378 at once. Contains diphacinone, an anticoagulant with a half-life in the dog of unknown days. For dogs that have ingested or that are suspected of having ingested diphacinone, and/or have obvious poisoning symptoms, such as bleeding or have elevated prothrombin times, give Vitamin K as follows: intramuscular injection. For anticoagulants with long half-lives, if known, it might be necessary to check prothrombin times every 3 days until values return to normal. See 'Note to Physician' for additional information.

California Department of Food and Agriculture  
1220 "N" Street, Room 341  
Sacramento, CA 95814

EPA SLN No. CA 890020  
EPA EST. No. XXXX-CA-X  
California Reg. No. 10965-50001

Net Contents:

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS PRODUCT MAY ONLY BE DISTRIBUTED AND USED IN CALIFORNIA AND MAY ONLY BE SOLD BY THE OFFICES OF CALIFORNIA'S COUNTY AGRICULTURAL COMMISSIONER

**READ THIS LABEL:** Read this entire label and follow all directions and use precautions.

**IMPORTANT:** Store product not in use at a location out of reach of children, pets, domestic animals, and wildlife. Dispose of product container, unused, spoiled, and unconsumed bait as specified in the "STORAGE AND DISPOSAL" section of this label.

#### USE RESTRICTIONS – FOR ALL USES

This product may only be used to control CALIFORNIA GROUND SQUIRRELS (*Spermophilus beecheyi*), NORWAY RATS (*Rattus norvegicus*), ROOF RATS (*R. rattus*), WOOD RATS (*Neotoma spp.*), MEADOW MICE (*Voles, Microtus californicus, M. montanus*), JACKRABBITS (Black-tailed hare, *Lepus californicus*), COTTONTAIL RABBITS (*Sylvilagus audobonii ssp.*), CHIPMUNKS (*Eutamias spp.*) and MUSKRATS (*Ondatra zibethica*) and at the use sites and using the application methods identified in the "USE DIRECTIONS" paragraphs indicated below for the target species groupings. Contact your local County Agricultural Commissioner's office if you need help in identifying the target species you intend to control or are not sure about the limits of site designations.

Use only in rodent infested areas where exposure to agricultural crops or commodities will not occur. **Do not** contaminate food or food stuffs.

**Do not** apply this product in or around homes or other human residences. **Do not** apply this bait at sites or to control pests not indicated on this label. **Do not** apply this product by application methods that are not specified on this label. **Do not** pile bait.

For certain use patterns, this product may be applied by use of spot baiting techniques as described below. For all other applications, bait must be used in covered or enclosed bait stations designed to minimize spillage and to limit or prevent access by nontarget species. Use bait stations which have entrances large enough to accommodate target species but not larger animals. Depending on the claimed pest being targeted, bait stations may include tamper-resistant designs, other types of covered or enclosed feeders or dispensers, covered nursery flats, or, for muskrats only, floating covered or enclosed bait stations. Where necessary to protect endangered rodents or where otherwise applicable, use the modified bait station designs described at <http://www.cdpr.ca.gov/docs/endspec/prescint.htm>. Secure bait stations to prevent overturning and bait spillage.

Apply baits in locations out of reach of children, pets, and domestic animals. If this is not possible, baits shall be used only in tamper-resistant bait stations that are resistant to destruction by dogs, wildlife, domestic animals, and children under six years of age. If bait is to be used in and around agricultural buildings and facilities for domestic animals the bait must be placed in tamper-resistant bait stations which shall be secured indoors or within 10 feet of farm buildings (other than farm or ranch house), fence lines, food or feed storage areas, lumber piles, or trash and garbage containers.

**Do not** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Keep all other persons out of the treatment area during application.

**Do not** place bait in areas where contaminating food, or surfaces that may contact food or feed, is likely.

**Do not** graze livestock or plant food or feed crops in spot-treated areas while bait is present. Applications in vineyards, orchards, and groves may only be made after harvest and during the dormant period and may not be made after tree and vine growth resumes in the spring.

### **BAIT STATION APPLICATIONS - GENERAL INSTRUCTIONS**

It may take several days or longer for target animals to become accustomed to a bait station and to begin to accept bait from it. Maintain an uninterrupted supply of bait in the bait stations for as long as target species are taking bait, which often will last from 1 to 4 weeks after feeding begins. Check stations one or more times per week and replace consumed, spoiled or contaminated bait. Properly dispose of bait that is removed from bait stations or is spilled or scattered from bait stations.

When toxic bait in bait station is no longer being accepted by target species, remove and dispose of it properly. Bait stations may be removed at the time or left in place if reinfestation is likely. Stations left in place may be baited periodically with nontoxic feed to aid in the detection of target species attempting to recolonize the site and to condition them to feeding in the bait stations. If target rodents return, census baits may be replaced by this product.

### **FOLLOW-UP OPERATIONS – FOR ALL USES**

Collect dead rodents and dispose of them by deep burying, burning (if permitted in your County or community), or double plastic bagging or by wrapping in newspaper and discarding in the trash. Wear disposable plastic gloves or other suitable hand protection if you must pick up carcasses by hand.

### **IN/AROUND AGRICULTURAL BUILDINGS, AGRICULTURAL CROPS, RANGELAND, FORESTRY PLANTATIONS AND SELECTED NONCROP AREAS**

#### **CALIFORNIA GROUND SQUIRRELS**

**USE RESTRICTIONS:** This product may be used to control California Ground Squirrels (*Spermophilus beecheyi*) in bait station applications in and around livestock buildings (e.g., cattle barns, poultry houses); around (but not within) livestock pens; in and around vineyards, orchards, and groves; in rangelands, noncrop borders, and fallow lands; along outsides of fence rows and rights-of-way adjacent to canal banks, ditch banks, highways, levees, railroad lines, and utilities; and in campgrounds, recreational areas, horticultural nurseries, and plantations of forest trees.

This product may also be used in spot-treatment applications in vineyards, orchards, and groves after harvest and during the dormant season, and in rangeland. Do not graze livestock or plant food or feed crops in spot-treated areas while bait is present.

**BAIT ACCEPTANCE TRIALS:** Before applying toxic bait, test for target species' readiness to accept this product by applying untreated crimped oat groats, to portions or all of the infested area, using the same application method (bait station or spot baiting) that is to be used for toxic baiting. Observe prebaited areas to determine whether nontarget species have found and consumed the oats. Adjust treatment methods to minimize nontarget hazards. Do not apply toxic bait if it appears that nontarget species are primarily responsible for consuming the oats. Apply toxic bait only if ground squirrels appear to have accepted the oats readily. Untreated crimped oat groats may be obtained from the County Agricultural Commissioner's office.

**BAIT STATION BAITING:** Secure tamper-resistant bait stations near active ground squirrel burrows or runways. Place stations at intervals of 20 to 100 feet. Load 1 to 5 pounds of bait into bait station. Inspect stations at least weekly and replenish bait as needed. Remove and properly dispose of spoiled or fouled bait.

**SPOT BAITING:** Using a bait spoon, evenly scatter 1/3 cup (0.1 lb) of bait over 40 to 50 square feet near active squirrel burrows and runways. Do not over-bait. Do not place bait in piles. Using the same procedure, make a second application 4 days after the first. Do not apply more than 10 pounds of bait per acre per treatment.

#### **NORWAY RATS and ROOF RATS**

**USE RESTRICTIONS:** This product may be used to control Norway rats (*Rattus norvegicus*) and roof rats (*R. Rattus*) in and around agricultural buildings and facilities for domestic animals (e.g., cattle barns, poultry houses); around (but not within) livestock pens; after harvest and in dormant season applications only in orchard, groves, and vineyards; rangeland, noncrop borders, and fallow lands; along outsides of fence rows and rights-of-way adjacent to canal banks, ditch banks, highways, levees, railroad lines, and utilities; and in campgrounds, recreational areas, horticultural nurseries, and plantations of forest trees.

**BAIT STATION BAITING:** Place 4 to 16 ounces of bait in a secure tamper-resistant bait station. Place bait stations in dry locations such as in concealed places, in corners, along walls where rats feed, drink, or frequent. For Norway rats, put bait stations at or near ground level and at burrows and harborages. For roof rats, place bait stations at ground floor and top floor, or attic levels. Inspect stations at least weekly and replenish bait as needed. Remove and properly dispose of spoiled or fouled bait.

#### **WOOD RATS**

**USE RESTRICTIONS:** This product may be used to control wood rats (*Neotoma spp.*) in and around agricultural buildings and facilities for domestic animals (e.g., cattle barns, poultry houses); after harvest and in dormant season applications only in orchard, groves, and vineyards; rangeland, noncrop borders, and fallow lands; along outsides of fence rows and rights-of-way adjacent to canal banks, ditch banks, highways, levees, railroad lines, and utilities; and in campgrounds, recreational areas, horticultural nurseries, and plantations of forest trees.

**BAIT STATION BAITING:** Place 4 to 16 ounces of bait in a secure tamper-resistant bait station. Bait stations should be located near existing wood rat runways or dens, and generally spaced no further than 100 feet apart. Inspect stations at least weekly and replenish bait as needed. Remove and properly dispose of spoiled or fouled bait.

### **AGRICULTURAL CROPS, RANGELAND, FORESTRY PLANTATIONS AND SELECTED NONCROP AREAS**

#### **MEADOW MICE (VOLES)**

**USE RESTRICTIONS:** This product may be used to control California voles and montane voles after harvest and in dormant season applications only in orchard, groves, and vineyards; in noncrop borders and fallow lands; along the outside of fence rows and rights-of-way adjacent to canal banks, ditch banks, highways, levees, railroad lines, and utilities; in campgrounds, recreational areas, horticultural nurseries and plantations of forest trees, and in rangelands. Do not graze livestock or plant food or feed crops in spot-treated areas while bait is present.

**SPOT BAITING:** Using a bait spoon, evenly scatter one to two tablespoons of bait (1/4 to 1/2 ounce) near active burrow openings or in runways. Do not over-bait. Do not place bait in piles. Using the same procedure, make a second application 4 days after the first. Do not apply more than 10 pounds of bait per acre per treatment.

#### **BLACK-TAILED JACKRABBITS (BLACK-TAILED HARES)**

**USE RESTRICTIONS:** This product may be used to control black-tailed jackrabbits (*Lepus californicus*) using self-dispensing feeders situated along borders of agricultural crops; in rangelands and fallow land; along outsides of fence rows and rights-of-way adjacent to canal banks, ditch banks, highways, levees, railroad lines, and utilities; in horticultural nurseries and plantations of forest trees; and in campgrounds and recreational areas.

**BAIT STATION BAITING:** Place 1 to 5 pounds of bait in a covered or enclosed self-dispensing feeder or a covered or enclosed nursery flat near runways, or nesting or feeding areas used by jackrabbits. Replenish bait as needed to control jackrabbits. Move feeders to new locations in permitted use sites if necessary to achieve acceptance of bait by jackrabbits. Remove and properly dispose of spoiled or fouled bait.

#### **COTTONTAIL RABBITS**

**USE RESTRICTIONS:** This product may be used to control cottontail rabbits (*Sylvilagus audobonii ssp.*) using self-dispensing feeders situated along borders of agricultural crops; in rangelands and fallow land; along outsides of fence rows and rights-of-way adjacent to canal banks, ditch banks, highways, levees, railroad lines, and utilities; in horticultural nurseries and plantations of forest trees; and in campgrounds and recreational areas.

**BAIT STATION BAITING:** Place 1 to 5 pounds of bait in a covered or enclosed self-dispensing feeder or a covered or enclosed nursery flat near runways, or nesting or feeding areas used by cottontails. Replenish bait as needed to control cottontails. Move feeders to new locations in permitted use sites if necessary to achieve acceptance of bait by cottontails. Remove and properly dispose of spoiled or fouled bait.

### **FORESTRY PLANTATIONS AND SELECTED NONCROP AREAS**

#### **CHIPMUNKS**

**USE RESTRICTIONS:** This product may be used to control chipmunks in tamper-resistant bait station applications in and around campgrounds and recreational areas; along noncrop borders; in fallow lands; along outsides of fence rows and rights-of-way adjacent to canal banks, ditch banks, highways, levees, railroad lines, and utilities; and in horticultural nurseries and plantations of forest trees.

**BAIT STATION BAITING:** Secure tamper-resistant bait stations near active chipmunk burrows or runways. Place stations at intervals of 20 to 50 feet. Load 4 to 16 ounces of bait into each bait station. Inspect stations at least weekly and replenish bait as needed. Remove and properly dispose of spoiled or fouled bait.

### **NATURAL/MANMADE WATERWAYS AND WETLANDS ADJACENT TO AGRICULTURAL CROPS, RANGELAND, FORESTRY PLANTATIONS AND SELECTED NONCROP AREAS**

#### **MUSKRATS**

**USE RESTRICTIONS:** This product may be used to control muskrats (*Ondatra zibethica*) in floating bait station applications in natural and manmade waterways and wetlands adjacent to agricultural crops, rangelands, noncrop borders, uncultivated agricultural areas and rights-of-way.

Do not apply baits to control muskrats in crops, rangeland, noncrop borders, uncultivated agricultural areas, and rights-of-way themselves

**BAIT STATION BAITING:** Secure covered or enclosed bait station to a small raft. Anchor or secure floating bait station to bank or bottom. Load 1 to 5 pounds of bait in a floating covered or enclosed bait station. Check stations at least weekly and replenish bait as needed to control muskrats. Remove and properly dispose of spoiled or fouled bait.

Revised 2-24-11

FIFRA SECTION 2(ee) RECOMMENDATION FOR USE  
IN THE STATE OF CALIFORNIA

**CDFA RODENT BAIT BLOCK DIPHACINONE**

EPA Reg. No. 61282-26-10965

Use Directions for Muskrats (*Ondatra zibethica*)

It is a violation of federal law to use this product in any manner inconsistent with its labeling. This recommendation is permitted under FIFRA Section 2(ee). This supplemental label and the federally registered label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on the EPA registered label.

**DIRECTIONS FOR USE**

**BAITING: MUSKRATS, (*Ondatra zibethica*)** Apply 1 to 16 CDFA Rodent Bait Blocks per bait station placement. Each CDFA Rodent Bait Block weighs approximately one ounce. Bait stations should be located near existing muskrat runways or dens. Floating bait stations may also be used. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of muskrat activity cease.

FIFRA SECTION 2(ee) RECOMMENDATION FOR USE  
IN THE STATE OF CALIFORNIA

**CDFA RODENT BAIT BLOCK DIPHACINONE**

EPA Reg. No. 61282-26-10965

Use Directions for Wood Rats (*Neotoma* spp.)

It is a violation of federal law to use this product in any manner inconsistent with its labeling. This recommendation is permitted under FIFRA Section 2(ee). This supplemental label and the federally registered label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on the EPA registered label.

**DIRECTIONS FOR USE**

**BAITING: WOOD RATS (*Neotoma* spp.)** Apply 1 to 16 CDFA Rodent Bait Blocks per bait station placement. Each CDFA Rodent Bait Block weighs approximately one ounce. Bait stations should be located near existing wood rat runways or dens, Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of wood rat activity cease.

## PRECAUTIONARY STATEMENTS

### Hazard to Humans and Domestic Animals

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

### Environmental Hazards

This product is toxic mammals and birds. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high-water mark.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**Storage:** Store only in original closed container in a cool, dry place inaccessible to children and pets. Store separately from fertilizer and away from products with strong odors which may contaminate the bait and reduce acceptability. Sweep up spillage carefully and collect for disposal.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** **Commercial Use:** Dispose of empty container in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **Household Use Disposal-** If empty: Do not reuse empty container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste disposal agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**NOTICE:** Read the entire Directions For Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be returned.

### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

Manufacturer and Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Manufacturer, and Buyer and User assume the risk of any such use. MANUFACTURER AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. THE EXCLUSIVE REMEDY OF USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES, OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF THE MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT. Manufacturer and Seller offer this product and Buyer and User accepts it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of manufacturer.



CALIFORNIA DEPARTMENT OF  
FOOD AND AGRICULTURE

## CDFA RODENT BAIT BLOCK DIPHACINONE

Mold and Moisture Resistant

Kills Norway rats, roof rats, and house mice in wet or dry areas

Active Ingredient:

Diphacinone [2-diphenylacetyl]-1,3-indandione].....0.005%

Inert Ingredients..... 99.995%

Total:.....100.000%

## KEEP OUT OF REACH OF CHILDREN CAUTION

### FIRST AID

Have label with you when obtaining treatment advice.

|                     |   |
|---------------------|---|
| <b>If Swallowed</b> | <ul style="list-style-type: none"><li>Call a poison control center, doctor or 1-800-498-5743 immediately for treatment advice.</li><li>Have person sip a glass of water if able to swallow.</li><li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li></ul> |
|---------------------|---|

### NOTE TO PHYSICIAN

If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally, as indicated in bis-hydroxycoumarin overdose. Repeat as necessary based on monitoring of prothrombin times.

Distributed by: California Department of Food and Agriculture  
1220 "N" Street, Room A-357  
Sacramento, CA 95814

EPA Reg. No. 61282-26-10965

EPA EST. No. XXXX-CA-X

Net Contents:

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all directions and use precautions. **IMPORTANT:** Do not expose children, pets, or other non-target animals to rodenticides. To help to prevent accidents:

1. Store product not in use at a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potential destructive animals, or in areas prone to vandalism.
3. Dispose of product container. And unused, spoiled, and unconsumed baits as specified on this label.

### USE RESTRICTIONS

For the control of Norway rats, roof rats, and house mice in and around homes, industrial and agricultural buildings, and similar man-made structures. This product is especially suited for use in wet or damp areas, including river banks, gullies, irrigation ditches, sewers, garbage dumps, and landfills.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come direct contact with food. Do not apply bait directly to ground surface or in grass or other ground cover.

### SELECTION OF TREATMENT AREAS

Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodent or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

### APPLICATION DIRECTIONS

Each CDFA Rodent Bait Block weighs approximately one ounce.

### RATS

Apply 4 to 16 CDFA Rodent Bait Blocks (usually at intervals of 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

### MICE

Place one 1-ounce CDFA Rodent Bait Block at each placement location. Space placements at 8 to 12 foot intervals. Two 1-ounce pieces may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

### RATS AND MICE

Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

Revised 11-18-04